

20 April 2026

Tozorakimab met primary endpoint in Phase III MIRANDA trial in patients with COPD

Third positive pivotal Phase III clinical trial of AstraZeneca's IL-33-targeting biologic further demonstrates its benefits in COPD

Positive high-level results from the pivotal Phase III MIRANDA trial showed potential first-in-class tozorakimab demonstrated a statistically significant and clinically meaningful reduction in the annualised rate of moderate-to-severe COPD exacerbations in the primary population of former smokers and in the overall population, which included former and current smokers, and patients across all blood eosinophil* counts and all stages of lung function severity.

In MIRANDA, patients received tozorakimab 300mg or placebo on top of standard of care once every two weeks.¹ The trial enrolled patients with COPD still experiencing moderate-to-severe exacerbations while on inhaled standard of care.¹ These results follow an [announcement](#) in March of the positive high-level results from the pivotal Phase III OBERON and TITANIA trials studying tozorakimab at a four-week dosing interval.

Frank Sciruba, MD, FCCP, Professor of Pulmonary and Critical Care Medicine, University of Pittsburgh, Chief Investigator of LUNA programme said: "These results add to the growing body of evidence that indicates tozorakimab delivered meaningful clinical benefits for COPD patients who urgently need new treatment options. Up to half of patients today still experience exacerbations even when taking standard-of-care inhaled therapies, putting them at risk of serious health consequences including hospitalisation and even death."

Sharon Barr, Executive Vice President, BioPharmaceuticals R&D, AstraZeneca, said: "These data further demonstrate tozorakimab's exciting potential as a first-in-class biologic with a truly differentiated mechanism of action that inhibits the signalling of the reduced and oxidised forms of IL-33 to address underlying drivers of COPD. We look forward to sharing the data with regulators and the scientific community as soon as possible."

Tozorakimab was generally well tolerated with a favourable safety profile consistent with previous trials. The data will be submitted to regulatory authorities and shared with the scientific community at an upcoming medical meeting.

Nearly 400 million people are diagnosed with COPD, a heterogenous and progressive disease and the 3rd leading cause of death globally.^{2,3} Even when on inhaled standard of care, more than 50% of patients experience exacerbations, putting them at an increased risk of cardiopulmonary events and mortality.⁴⁻⁷

Tozorakimab is a potential first-in-class monoclonal antibody targeting interleukin-33 (IL-33), that uniquely inhibits the signalling of the reduced and oxidised forms of IL-33, offering the potential to both reduce inflammation and disrupt the cycle of mucus dysfunction that contribute to COPD worsening.⁸⁻¹¹

Tozorakimab is also being studied in a Phase III trial for severe viral lower respiratory tract disease and in a Phase II trial in asthma.^{12,13}

**eosinophil: a type of white blood cell, which at increased levels may contribute to inflammation in respiratory diseases.¹⁴*

Notes

COPD

COPD, the third leading cause of death (excluding COVID-19) worldwide, is a progressive respiratory condition characterised by persistent airflow limitation and chronic inflammation of the airways.^{3,15} Common symptoms include breathlessness, chronic cough and excess mucus production.¹⁵ These symptoms can worsen over time and contribute to ongoing inflammation and bronchoconstriction, making it difficult to breathe and increasing the risk of COPD exacerbations.¹⁵ These COPD exacerbations have a profound impact on the lives of those with the

disease, accelerating disease progression, increasing hospitalisations, and increasing the risk of future cardiopulmonary events - including heart attacks, all of which can be life-threatening.^{7,15} In the US, exacerbations cause more than 2,500 emergency department visits per day.¹⁶ Only 50% of COPD patients live more than 3.5 years after their first severe exacerbation.¹⁷

Phase III LUNA programme

Tozorakimab's Phase III COPD development programme includes four clinical trials: OBERON, TITANIA, MIRANDA and PROSPERO.

OBERON and TITANIA

OBERON and TITANIA are replicate Phase III double-blind, placebo-controlled trials investigating the efficacy and safety of tozorakimab in adults with symptomatic COPD with a history of ≥ 2 moderate or ≥ 1 severe COPD exacerbations in the 12 months prior to enrolment. A total of 2,306 patients were randomised in both trials, including former and current smokers, and patients across all blood eosinophil counts and all stages of lung function severity.^{18,19} Patients were placed on a regimen of tozorakimab 300mg once every four weeks, or placebo over the course of 52 weeks on top of inhaled therapy.

Prior to enrolment, patients received standard of care inhaled maintenance therapy for at least three months. The primary endpoint is annualised rate of moderate-to-severe COPD exacerbations in former smokers with COPD. A key secondary endpoint measured the annualised rate of moderate-to-severe COPD exacerbations in the overall population of former and current smokers.^{18,19}

MIRANDA

MIRANDA is a Phase III double-blind, placebo-controlled trial investigating the efficacy and safety of tozorakimab in adults with symptomatic COPD with a history of ≥ 2 moderate or ≥ 1 severe COPD exacerbations in the 12 months prior to enrolment. A total of 1,454 patients were randomised in this trial, including former and current smokers, and patients across all blood eosinophil counts and all stages of lung function severity.¹ Patients were placed on a regimen of tozorakimab 300mg once every two weeks, or placebo over the course of 52 weeks on top of inhaled therapy.

Prior to enrolment, patients received standard of care inhaled maintenance therapy for at least three months. The primary endpoint is annualised rate of moderate-to-severe COPD exacerbations in former smokers with COPD. A key secondary endpoint measured the annualised rate of moderate-to-severe COPD exacerbations in the overall population of former and current smokers.¹

PROSPERO

The PROSPERO trial is a randomised, long-term extension clinical trial that enrolled patients who completed the OBERON or TITANIA trials. PROSPERO has a different primary endpoint, which is the annualised rate of only severe COPD exacerbations (hospitalisations and death) in former smokers with COPD over 104 weeks. A total of 1,713 patients were randomised in this trial.²⁰

Tozorakimab

Tozorakimab is being developed by AstraZeneca as a first-in-class potent human immunoglobulin monoclonal antibody that binds to interleukin (IL-33). Tozorakimab targets the top of the inflammatory cascade uniquely inhibiting IL-33 signalling in two ways, thereby suppressing inflammation and disrupting the cycle of mucus dysfunction.⁸ Tozorakimab is currently being investigated in a Phase III clinical trial for severe viral lower respiratory tract disease and a Phase II trial for asthma.^{12,13} Tozorakimab was granted Fast Track Designation by the US Food and Drug Administration for the treatment of severe viral lower respiratory tract disease in November 2023 and for COPD in December 2024.²¹

AstraZeneca in Respiratory & Immunology

Respiratory & Immunology, part of AstraZeneca BioPharmaceuticals, is a key disease area and growth driver to the Company.

AstraZeneca is an established leader in respiratory care with a 50-year heritage and a growing portfolio of medicines in immune-mediated diseases. The Company is committed to addressing the vast unmet needs of these chronic, often debilitating, diseases with a pipeline and portfolio of inhaled medicines, biologics and new modalities aimed at previously unreachable biologic targets. Our ambition is to deliver

life-changing medicines that help eliminate COPD as a leading cause of death, eliminate asthma attacks and achieve clinical remission in immune-mediated diseases.

[AstraZeneca](#)

AstraZeneca (LSE/STO/NYSE: AZN) is a global, science-led biopharmaceutical company that focuses on the discovery, development, and commercialisation of prescription medicines in Oncology, Rare Diseases, and BioPharmaceuticals, including Cardiovascular, Renal & Metabolism, and Respiratory & Immunology. Based in Cambridge, UK, AstraZeneca's innovative medicines are sold in more than 125 countries and used by millions of patients worldwide. Please visit astrazeneca.com and follow the Company on Social Media [@AstraZeneca](#).

Contacts

For details on how to contact the Investor Relations Team, please click [here](#). For Media contacts, click [here](#).

References

1. Clinicaltrials.gov. Efficacy and Safety of Tozorakimab in Symptomatic Chronic Obstructive Pulmonary Disease With a History of Exacerbations (MIRANDA). [Online]. Available at: <https://clinicaltrials.gov/study/NCT06040086>. [Last accessed: April 2026]
2. Montes de Oca M, Perez-Padilla R, Celli B, et al. The global burden of COPD: epidemiology and effect of prevention strategies. *Global Epidemiology of Chronic Respiratory Disease*. 2025; 13(18):709-724.
3. World Health Organization (WHO). The top 10 causes of death. 2024. Available at: <https://www.who.int/news-room/fact-sheets/detail/the-top-10-causes-of-death>. [Last accessed: April 2026]
4. Chen S, Small M, Lindner L, Xu X. Symptomatic burden of COPD for patients receiving dual or triple therapy. *Int J Chron Obstruct Pulmon Dis* 2018;13:1365-1376.
5. Chen S, Miravittles M, Rhee CK, et al. Patients with chronic obstructive pulmonary disease and evidence of eosinophilic inflammation experience exacerbations despite receiving maximal inhaled maintenance therapy. *Int J Chron Obstruct Pulmon Dis* 2022;17:2187-2200.
6. Nordon C, Carstens D, Fageras M, et al. Exacerbation and mortality in COPD patients on triple inhaler and at high exacerbation risk. *Eur Respir J* 2024;64(Suppl. 68):PA1287 (Abstract).
7. Singh D, Han MK, Hawkins NM, et al. Implications of cardiopulmonary risk for the management of COPD: a narrative review. *Adv Ther*. 2024;41:2151-2167.
8. England E, Rees DG, Scott IC, et al. Tozorakimab (MEDI3506): an anti-IL-33 antibody that inhibits IL-33 signalling via ST2 and RAGE/EGFR to reduce inflammation and epithelial dysfunction. *Sci Rep*. 2023;13:9825.
9. Singh D, Guller P, Reid F, et al. A Phase 2a trial of the IL-33 mAb tozorakimab in patients with COPD: FRONTIER-4. *Eur Respir J* 2025;doi 10.1183/13993003.02231-2024.
10. Strickson S, Houslay KF, Negri VA, et al. Oxidised IL-33 drives COPD epithelial pathogenesis via ST2-independent RAGE/EGFR signalling complex. *Eur Respir J* 2023;62:2202210.
11. Strickson S, et al. Distinct Pharmacological Profiles of IL-33 Antibodies. [Poster Presentation]. Presented at the American Thoracic Society (ATS) 2024. May 2024.
12. Clinicaltrials.gov. Efficacy and Safety of Tozorakimab in Patients Hospitalised for Viral Lung Infection Requiring Supplemental Oxygen (TILIA). [Online]. Available at: <https://clinicaltrials.gov/study/NCT05624450> [Last accessed: April 2026]
13. Clinicaltrials.gov Dose Range Finding Study to Assess Efficacy and Safety of Tozorakimab in Adults With Uncontrolled Asthma on Medium-to-High Dose Inhaled Corticosteroids (UMBRIEL). [Online]. Available at: <https://clinicaltrials.gov/study/NCT06932263> [Last accessed: April 2026]
14. Jackson D, Akuthota P, Roufosse F. Eosinophils and eosinophilic immune dysfunction in health and disease. *Eur Respir Rev* 2022 31(163):210150.
15. GOLD. Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Pulmonary Disease: 2026 Report. Available at: <https://goldcopd.org/2026-gold-report-and-pocket-guide/> [Last accessed: April 2026]
16. American Lung Association. COPD Trends Brief: Burden. Available at: <https://www.lung.org/research/trends-in-lung-disease/copd-trends-brief/copd-burden> [Last accessed: April 2026]

17. Suissa S, Dell'Aniello S, Ernst P. Long-term natural history of chronic obstructive pulmonary disease: severe exacerbations and mortality. *Thorax* 2012;67:957-963.
18. Clinicaltrials.gov. Efficacy and Safety of Tozorakimab in Symptomatic Chronic Obstructive Pulmonary Disease With a History of Exacerbations (OBERON). Available at: <https://clinicaltrials.gov/study/NCT05166889>. [Last accessed: April 2026]
19. Clinicaltrials.gov. Efficacy and Safety of Tozorakimab in Symptomatic Chronic Obstructive Pulmonary Disease With a History of Exacerbations. (TITANIA). [Online]. Available at: <https://clinicaltrials.gov/study/NCT05158387>. [Last accessed: April 2026]
20. Clinicaltrials.gov. Long-term Efficacy and Safety of Tozorakimab in Participants With Chronic Obstructive Pulmonary Disease With a History of Exacerbations (PROSPERO). [Online]. Available at: <https://clinicaltrials.gov/study/NCT05742802> [Last accessed: April 2026]
21. AstraZeneca. Clinical Trials Appendix Q3 2025 Results Update. Available from: <https://www.astrazeneca.com/content/dam/az/PDF/2025/9m-q3/9M-and-Q3-2025-results-clinical-trials-appendix.pdf> [Last accessed: April 2026]

Matthew Bowden
Company Secretary
AstraZeneca PLC

This information is provided by RNS, the news service of the London Stock Exchange. RNS is approved by the Financial Conduct Authority to act as a Primary Information Provider in the United Kingdom. Terms and conditions relating to the use and distribution of this information may apply. For further information, please contact rns@lseg.com or visit www.rns.com.